

# Biology and genetic engineering of fruit maturation for enhanced quality and shelf-life

Antonio J Matas<sup>1,\*</sup>, Nigel E Gapper<sup>2,\*</sup>, Mi-Young Chung<sup>2</sup>,  
James J Giovannoni<sup>2,3</sup> and Jocelyn KC Rose<sup>1</sup>

Commercial regulation of ripening is currently achieved through early harvest, by controlling the postharvest storage atmosphere and genetic selection for slow or late ripening varieties. Although these approaches are often effective, they are not universally applicable and often result in acceptable, but poor quality, products. With increased understanding of the molecular biology underlying ripening and the advent of genetic engineering technologies, researchers have pursued new strategies to address problems in fruit shelf-life and quality. These have been guided by recent insights into mechanisms by which ethylene and a complex network of transcription factors regulate ripening, and by an increased appreciation of factors that contribute to shelf-life, such as the fruit cuticle.

## Addresses

<sup>1</sup> Department of Plant Biology, Cornell University, Ithaca, NY 14853, USA

<sup>2</sup> Boyce Thompson Institute for Plant Research, Cornell University, Ithaca, NY 14853, USA

<sup>3</sup> US Department of Agriculture – Agricultural Research Service, Robert Holly Center for Agriculture and Health, Ithaca, NY 14853, USA

\* These authors contributed equally to this manuscript and should be considered co-first authors.

Corresponding author: Rose, Jocelyn KC ([jr286@cornell.edu](mailto:jr286@cornell.edu))

Current Opinion in Biotechnology 2009, 20:197–203

This review comes from a themed issue on  
Food biotechnology  
Edited by Max Teplitski and Anita Wright

Available online 30th March 2009

0958-1669/\$ – see front matter

© 2009 Elsevier Ltd. All rights reserved.

DOI [10.1016/j.copbio.2009.02.015](https://doi.org/10.1016/j.copbio.2009.02.015)

Fruits are important contributors to human diets and health, providing essential nutrients, antioxidants, carbohydrates, and fiber. The ripening process has evolved to make fruit palatable to organisms that consume them and disperse their seeds. In doing so, ripening activates pathways that generally influence the levels of pigments (typically carotenoids and flavonoids), sugars, acids, and aroma volatiles, to make the organ more appealing, while simultaneously promoting tissue softening and degradation to permit easier seed release [1,2]. Indeed the volatiles produced by ripening fruit are also derived from, and represent signals for, the presence of essential nutrients for animals which may consume them [3]. Increased

susceptibility to postharvest microbial infection helps insure that the seed are released via fruit rot, if not consumption. A major challenge to breeders and producers of fruit species continues to be how to capture and deliver to market the desirable flavor, color, and texture attributes of ripe fruit while inhibiting or delaying ripening sufficiently to counter the negative consequences of over-ripening: oversoftening and decay.

## Ethylene and fruit ripening

Ripening is regulated by both internal and external stimuli, including temperature, light, plant nutrient status, water availability, and hormones. Many fruits, in particular the so-called climacteric fruits, require ethylene for ripening, resulting in the targeting of this hormone as a means of ripening control. Although nonclimacteric fruits typically produce little ethylene during ripening, many have still been shown to be affected by exogenous ethylene during ripening, making ethylene control a target for shelf-life manipulation even in species whose fruit do not require ethylene to ripen ([1,2] and references therein). Because of the importance of ethylene for ripening, the genes responsible for its synthesis, ACC synthase (ACS) and ACC oxidase (ACO), were early targets of study and manipulation and repression of using antisense strategies delayed ripening in either tomato or other species (reviewed in [2]). Commercialization of these technologies has been limited by the cost of regulatory compliance, real or perceived consumer concerns, and the efficiency of available postharvest ethylene control systems.

## Ethylene response genes: potential targets for ethylene response and ripening control

Ethylene signal transduction genes have been well characterized in *Arabidopsis* (for recent reviews see [4–6]). More recent efforts in understanding the ethylene response during fruit ripening have focused on the characterization of tomato homologs. All the components of the *Arabidopsis* ethylene signal transduction analyzed thus far are conserved in tomato (reviewed in [2]). However, the family size and expression profiles of some of these genes differ between *Arabidopsis* and tomato. For example, there are six ethylene receptors in tomato, compared to five in *Arabidopsis*, but all have similar binding affinities for ethylene. Further, expression studies indicate distinct profiles for *Arabidopsis* and tomato ethylene receptor genes. In *Arabidopsis* the *CONSTITUTIVE TRIPLE*

*RESPONSE1* (*CTR1*) MAP kinase kinase kinase is represented by a single locus, whereas in tomato a family of three genes exists, all of which can complement the *Arabidopsis ctr-1* loss-of-function mutation. The additional components of ethylene signal transduction in tomato point to more opportunities for fine-tuning regulation of the hormone response than in *Arabidopsis*. Recently, an exploration of the protein–protein interactions of tomato ethylene receptors and the downstream kinases revealed that a single ethylene receptor, *NEVER RIPE* (*NR*), is capable of interacting with multiple LeCTR proteins [7], supporting the hypothesis that ethylene receptors transmit the signal to downstream CTRs. However, it is still unclear whether members of the tomato CTR family are functionally redundant or elicit unique ethylene responses.

Recently, novel ethylene signaling components were discovered by the simultaneous cloning of the tomato *GREEN RIPE* (*GR*) and *Arabidopsis* REVERSION TO ETHYLENE SENSITIVITY1 (*RTE1*) loci. The gene responsible for the dominant ripening mutation in tomato was positionally cloned [8], whereas *RTE1* was discovered through a second-site suppressor screen of an *Arabidopsis* ETHYLENE RECEPTOR1 (*ETR1*) mutant [9]. There are at least two other GR-like genes in the tomato genome (*GRL1* and *GRL2*), but only one other family member of *RTE1*, designated *RTH*, is present in *Arabidopsis*. Characterization of the *GR* mutation revealed a deletion of noncoding 5'-upstream sequence that results in fruit-specific ectopic expression of this gene during early development and throughout ripening. This overexpression causes a severe reduction in ethylene sensitivity, which in turn results in drastic ripening inhibition even though normal or elevated ethylene levels are produced [10]. Constitutive overexpression of *GR* in transgenic plants caused inhibition of ripening but surprisingly little, if any, altered ethylene sensitivity was observed for other aspects of plant development. This suggests *GR* might interact with components of the fruit-specific ethylene response. Furthermore, the fact that the ethylene response inhibition results from ectopic expression, rather than targeted gene repression, suggests that the tomato gene might be functional in other species merely through overexpression.

The gene underlying the original ethylene-insensitive *Etr1* ethylene receptor mutation was isolated from *Arabidopsis* and shown to encode a two-component receptor kinase that binds ethylene (reviewed in [1,2]). The mutation in this and many other dominant ethylene-insensitive receptor alleles impaired the ability of the receptor protein to bind ethylene. Thus, another strategy for blocking ethylene perception in fruit could be at the receptor itself. Indeed, overexpression of the mutant *ETR1* protein resulted in delayed fruit ripening in tomato and inhibition of petal senescence in petunia. The tomato

*Never-ripe* (*Nr*) mutation was also shown to encode an ERS-like ethylene receptor that is also impaired in the ability to bind ethylene. Transcript levels of *LeETR1*, *LeETR2*, and *LeETR5* change little upon treatment of ethylene in fruit, whereas *NR*, *LeETR4*, and *LeETR6* increase during ripening, suggesting that they may be more important in fruit [11]. Reduction of either of two ethylene receptors, *LeETR4* or *LeETR6*, causes an early ripening phenotype in tomato, in agreement with models where ethylene receptors act through kinase-mediated inhibition of the negative regulator CTR1 [11]. Further, the receptor proteins are targets of the 26S proteasome-dependent pathway, and are degraded in the presence of ethylene. In addition, exposure of immature fruits to ethylene caused a reduction in the amount of ethylene receptor protein and earlier ripening, supporting the hypothesis that receptor levels regulate the timing of the onset of fruit ripening by measuring cumulative ethylene exposure [11]. Recently a tetratricopeptide repeat protein (*SITPR1*) was isolated from tomato and shown to interact in yeast two hybrid assays with *NR* and *LeETR1* proteins [12]. This class of protein is involved in the ubiquitination process which leads to proteasome-mediated protein degradation. Ethylene and auxin responses were modulated in plants overexpressing *SITPR1*; however, fruit ripening, as monitored by color and textural changes, was not affected. It is proposed that a similar and specific protein–protein interaction occurs for *LeETR4* and *LeETR6* to regulate receptor levels during tomato fruit ripening. In summary, the receptors, and the proteins with which they interact, provide additional targets for the manipulation of ethylene responses, including ripening. The broad array of responses regulated by ethylene suggests that for such genes to be effective in ripening control they would need to be modified in a fruit-specific manner.

### Transcriptional control of fruit ripening and conserved ripening regulators

Although considerable effort has focused on ethylene synthesis and response, only recently have inroads been made into understanding ripening control before ethylene: a regulatory system that is possibly conserved between climacteric and nonclimacteric species. Recently, the mutated genes underlying the ripening mutants *RIPENING INHIBITOR* (*RIN*) and *COLOURLESS NON RIPENING* (*CNR*) were cloned, providing key insights into the regulation of the ripening process upstream of ethylene. The *rln* and *Cnr* mutations are recessive and dominant mutations, respectively, which block the ripening process and result in fruit that are unable to respond to ripening-associated ethylene [13,14]. The *rln* mutation is especially interesting as it has been bred into many commercial tomato varieties to slow ripening and extend shelf-life. Both *rln* and *Cnr* encode transcription factors, providing insights into dedicated fruit-specific control of ripening. *rln* encodes a

MADS-box protein of the SEPALLATA clade [15], while *Cnr* is an epigenetic mutation that alters the promoter methylation of a SQUAMOSA promoter binding (SBP) protein. Both these transcription factors are required for ripening-associated increases in respiration and ethylene; characteristics of tomato and other climacteric fruit species. Furthermore, fruits from both mutants remain responsive to ethylene (even though they do not ripen), in that ethylene-responsive genes are induced by exogenous ethylene. The fact that they do not ripen in response to the hormone indicates that both genes operate upstream of crucial ripening activities, including ethylene biosynthesis and necessary activities that are apparently ethylene independent (Figure 1). This observed ethylene-independent aspect of ripening physiology suggests that both RIN-MADS and CNR-SBP proteins could be conserved ripening regulators of fruit in both the climacteric and nonclimacteric categories. Strawberry is the most widely studied species of the nonclimacteric classification. A ripening-related strawberry homolog of the tomato *RIN* gene has been isolated, which is consistent with the hypothesis that such regulators might be conserved between both climacteric and nonclimacteric fruits [13]. *RIN* and *CNR* may therefore represent good candidates for ripening control in a broader array of species than is currently possible via ethylene control alone.


Another tomato transcription factor, *LeHB-1*, has also recently been characterized and shown to play a central

role in fruit ripening [16<sup>••</sup>]. *LeHB-1* encodes a putative HD-Zip homeobox protein that binds to the promoter of ACC oxidase (*LeACO1*), which encodes a key enzyme in the ethylene biosynthetic pathway. Antisense inhibition of *LeACO1* expression was reported to result in reduced ethylene synthesis and in this recent study, transient silencing of *LeHB-1* resulted in a significant delay of ripening and a reduction of *LeACO1* transcript abundance [16<sup>••</sup>]. Together these observations support the idea that *LeHB-1* positively regulates *LeACO1* and that silencing of *LeHB-1* represses *LeACO1*, consequently leading to a delay in ripening. Putative *LeHB-1*-binding sites are also present in a number of other ripening-related genes including *LeACO2*, *LePG1*, and *LeMADS-RIN* and so it is possible that *LeHB-1* might also regulate these ripening-related genes directly (Figure 1).

### Genetic manipulation of ripening regulatory genes

To date, attempts to manipulate the expression of target genes during fruit ripening have produced varying results. As mentioned, transcriptional control has been utilized with respect to breeding for heterozygosity at the *rin* mutation (*Rin/rin*) in commercial lines to delay ripening and extend shelf-life; however, this modification leads to the inhibition of flavor and nutritional compound accumulation along with undesirable textural traits in some backgrounds due to incomplete ripening. In addition, constitutive expression or silencing of other

Figure 1



Model for the molecular regulation in tomato ripening. This model was updated from [2] with *LeHB-1* elucidated by [16<sup>••</sup>]. The promoters of *LeACO1*, *LeACO2*, *LePG1*, and *LeMADS-RIN* contain a putative *LeHB-1*-binding site suggesting the *LeHB-1* transcription factor may universally control fruit ripening as well as floral organogenesis and carpel development.

target genes can cause unwanted pleiotropic effects on other aspects of plant development, or even lead to lethality as is probably the case for the *Cnr* and *LeHB-1* loci, as stable transgenic plants harboring either of these genes have been difficult to produce. As a consequence of unwanted pleiotropic effects, targeted analysis of the transgene effect in the fruit is often difficult, as exemplified by the light signal transduction and carotenoid regulatory gene *HIGH-PIGMENT2*, where pleiotropic effects of transgene expression in nonfruit tissues were noted (low yield and brittleness), although this can be overcome through the use of a fruit-specific promoter [17]. More accurate phenotypic determination of fruit function and utility for the assessment of other production or quality influences will be achieved through the use of appropriate tissue-specific and development-specific promoters. Indeed, the most useful transgenic or natural alleles for ripening control will be those that deliver fruit specificity and result in levels of expression sufficient for extended shelf-life while ripening sufficiently for desirable quality. While several fruit-specific promoters are known, the commercial use of most is subject to intellectual property restrictions and even they represent a narrow range of expression options (essentially either throughout fruit development or induced at the onset of ripening). Isolation and utilization of promoters with more desirable temporal or spatial expression profiles could provide the means to regulate fruit ripening by manipulating the expression of transcription factors such as *rin*, *Cnr*, and *LeHB-1*, absent unintended effects. Further, more detailed characterization of the molecular make up of ripening regulatory networks [18] should provide additional fruit-specific loci which can be altered genetically to avoid undesirable secondary effects, without the need of fruit-specific promoters. Some examples with particular reference to altering cell wall metabolism and texture are explored in subsequent sections.

### Cell walls and cuticles as key factors for fruit shelf-life

The cell wall has a profoundly important influence on fruit texture and cell wall components and the underlying genes have been frequent targets for genetic engineering, mostly in tomato, with the goal of extending shelf-life [19,20]. Although knowledge of the mechanism of fruit softening has grown in recent years, it has proven especially difficult to establish the relationship between specific aspects of cell wall metabolism or architecture and their relationship to changes in tissue firmness [19,21]. Similarly, it has also been difficult to establish the fine regulation of genetic elements that produce a range of effects depending on the genetic background in which they are introduced [22]. The regulation of texture and shelf-life is clearly far more complex than was previously envisaged and so new approaches are needed, including the inclusion of observations in species beyond the

traditional tomato experimental model [23]; a better understanding of the relationship between changes in the textural properties of specific fruit tissues, intact fruit 'firmness' and shelf-life; and more comprehensive models of the biochemical and physiological elements that contribute to fruit 'firmness'.

Ripening-related disassembly of the cell wall polysaccharide matrix is generally the only factor that is mentioned when describing the structural basis of fruit softening and the associated loss of shelf-life and fruit quality. This is largely a reflection of the relative ease with which some of the dramatic changes in wall polymer structure and composition, and the activities of the associated enzymes, can be measured. However, recent studies have started to examine the potential involvement and relative importance of other structures and physiological processes. For example, it has been proposed that differences in tomato fruit cuticle structure and composition may be associated with the substantial variation in tomato fruit shelf-life that has been reported in different tomato genotypes [24<sup>•</sup>]. The cuticle has a number of biological functions that could have an important impact on fruit quality and shelf-life, including the ability to maintain fruit skin integrity [25], restrict cuticular transpiration [26<sup>•</sup>], and limit microbial infection. Other reports have also highlighted ripening-related processes that likely contribute to fruit firmness, such as turgor pressure [24<sup>•</sup>,27<sup>•</sup>] and the possibly associated developmental changes in apoplastic solute accumulation [28]. Integration of all these features into a more holistic model of ripening-related textural changes should provide a host of new targets for gene manipulation, with the ultimate goal of allowing more complete fruit ripening on the plant, thereby promoting nutrient and flavor accumulation while maintaining fruit firmness and shelf-life.

### When a fruit is more than a fruit?

Fresh fruits and their processed derivatives represent important sources of carotenoids, flavonoids, and anthocyanins and the possibility of improving content using bioengineering represent an opportunity to improve access to healthy foods. The inherent resiliency of plant metabolism to maintain homeostasis has impaired efforts to facilitate changes in some of these pathways, as has been especially well documented for carotenoids [29]. An alternate approach is to focus on regulatory rather than biosynthetic genes. Accumulation of carotenoids in high-pigment fruit mutants has been reported to result from an increase in the plastid number, larger plastid compartment size, increased plastid division, and a greater capacity for pigment storage [30,31]. These changes in pigmentation, plastid type, and metabolism were associated with the elevation of transcripts from key carotenoid genes such as phytoene synthetase-1 (*Psy-1*) and are not directly connected with the ripening process [32], thus



allowing modifications in carotenoid content without affecting other aspects of fruit ripening or broader plant development. Moreover, this particular modification has been reported to be stable in field tests for transgene stability and yield performance [33].

Recently, a MYB transcription factor from snapdragon was expressed using the E8 promoter specifically in tomato fruit, causing elevated levels of anthocyanins: compounds that have been shown to increase the longevity of cancer-susceptible mice [34<sup>••</sup>]. A homolog of this transcription factor (MYB10) was previously cloned and characterized from apple and was shown to positively regulate the anthocyanin pathway [35]. Further, an allelic rearrangement in the promoter of MYB10 causes a novel autoregulatory motif, which is sufficient to account for the increase in MYB10 levels and subsequent ectopic accumulation of anthocyanins throughout the plant [36]. This modification was found in all red apple species tested, but not in white apple varieties. This gain-of-function mutation in the anthocyanin regulatory pathway has significant implications for the development of novel varieties of plants and fruit with enhanced nutritional status and increased consumer appeal, by using either genetic modification or by more conventional breeding methods.

Examples of specific pathways that have been targeted through genetic engineering with either enhanced shelf-life or nutritional status, or genotypes of particular interest are shown in Table 1. However, impacts on health and nutrition can be mediated through less direct means than the alteration of nutrition associated biochemical pathways. These goals can be met by modifying overall appeal and quality so that fruit and fruit products become more competitive choices for consumers and net fruit consumption is increased. For example, seedless tomato fruit were developed by silencing a key enzyme of the flavonoid biosynthesis pathway [37].

## New tools, new strategies, and enhancement of traditional approaches

Prior efforts toward fruit quality improvement have focused on single steps in biochemical pathways, but complex feedback regulation networks have complicated such strategies, as noted above. Targeting regulatory genes may provide a route to avoid these regulatory impediments though the difficult work of assessing the impact of altering specific pathway steps is required to understand the underlying regulatory networks [32]. The use of a transcription factor that regulates a complete biosynthesis pathway, as opposed to targeting a single enzyme, has proved useful to study and manipulate the accumulation of tannins in grape [38] or polyphenolic antioxidants in tomato [39]. New high-throughput functional genomics and systems approaches are also allowing more accurate predictions of regulatory processes [40]. Applying such methodologies to mapping of quantitative trait loci (QTL) allowed for the detection of epistatic interactions among genes [41<sup>•</sup>], the discovery of new genes related with metabolic pathways of interest, and even testing of the stability of QTL effects over multiple seasons and in different environments [42].

Recent reports revealed that the screening for genes whose expression varies between lines that are genetically very similar, and yet varying significantly in fruit quality, is a helpful tool for identifying potentially valuable genetic targets [43], and provides insights concerning the nature of complex genetic control processes that relate to fruit nutrient content [44], ripening, and softening [45]. These tools make it possible to identify potential candidate genes in model plants, like *Arabidopsis*, which could lead to targeting important genes in crop species [46]. In one interesting example, tomatoes were biofortified with calcium by the overexpression of an *Arabidopsis* calcium transporter (*CAX1*) [47]. However, while the calcium levels were increased, the bioavailability of the increased calcium was questioned. Similar experiments were carried out,

**Table 1**

**Recent transgenic and mutant tomato and grape genotypes and their function/effects associated with nutrition or shelf-life**

	Gene or mutant name/ID <sup>a</sup>	Biological process	Refs
<b>Shelf-life</b>			
β-Ketoacyl-CoA synthase	<i>LeCER6/3760026</i>	VLFC-cuticular waxes	[26 <sup>•</sup> ]
Cuticular water permeability	<i>Cwp1/100136893</i>	Cuticle microfissuring and dehydration	[25]
Delayed fruit deterioration	<i>dfd</i>	Extended shelf-life, reduced water loss	[24 <sup>•</sup> ]
Ethylene receptor	<i>LeETR4/543588</i>	Early-ripening fruit	[53]
Auxin response factor	<i>DR12</i>	Modification of fine pectin structure and tissue architecture	[21]
<b>Nutrient content</b>			
Chalcone synthase	<i>Chs/778294, 778295</i>	Flavonoid content reduction, parthenocarpic fruit	[37]
Phytoene synthase-1	<i>PSY-1/543988</i>	Changes to pigmentation, plastid type, and metabolism associated with Psy-1 overexpression	[32]
Isoflavone synthase	<i>GmIFS2/606705</i>	Presence of genistin in leaves	[54]
MYB transcription factor	<i>VvMYBPA1/100232899</i>	Specific regulation of proanthocyanidin biosynthesis	[38]
Flavonol-specific transcriptional activator	<i>AtMYB12/819359</i>	Flavonol accumulation in fruits	[39]

<sup>a</sup> Gene ID from [www.ncbi.org](http://www.ncbi.org).

where the intake of calcium from transgenic carrots over-expressing a *CAX1* gene were monitored in both mice and humans [48]. Calcium absorption in bones increased for both mice and humans eating *CAX1*-expressing carrots compared to controls, demonstrating alternative means of fortifying fruits and vegetables with bioavailable calcium.

## Prospects for the future

It is important to note that recent studies have reported that genetically modified organisms are not inherently dangerous and can be considered safe to consumers when compared with conventional foods developed for target traits through naturally occurring genetic variation [49]. Of course, given that the safety of crops is a primary concern for both producers and consumers, it is necessary to define the risk and unintended effects in genetically modified plants [50], including the effects of a specific gene and the possible effects in a susceptible subgroup of consumers [51]. Although many of the target genes that result from emerging genomics approaches may not in themselves be candidates for direct manipulation, they may represent DNA sequences that can be exploited as markers in breeding programs to select for allelic variation, which can be tested and potentially used for the alteration of target traits. Selection of QTLs in particular can be made much more efficient through marked-assisted selection (MAS). Integrating MAS in traditional breeding practices is most likely the highest short-term impact of our rapidly increasing understanding of fruit molecular biology, as it is economically attractive and applicable for markets in both developed and developing countries [52]. The current proliferation of genetically modified cereal and fiber crops is likely to alleviate consumer concerns, reduce the current regulatory costs, and result in more direct biotechnology applications to fruit crops in coming years.

## Acknowledgements

We acknowledge the support from USDA-NRI grants 2007-02773 and 2006-35304-17323, and the NSF Plant Genome grants 05-01778 and 06-06595. Support was also provided by a grant to JKCR from the US-Israel Binational Science Foundation, and to NEG from the New Zealand Foundation for Research, Science and Technology FRST-BTPR0701.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Giovannoni JJ: **Genetic regulation of fruit development and ripening.** *Plant Cell* 2004, **16**:S170-S180.
  2. Giovannoni JJ: **Fruit ripening mutants yield insights into ripening control.** *Curr Opin Plant Biol* 2007, **10**:283-289.
  3. Goff SA, Klee HJ: **Plant volatile compounds: sensory cues for health and nutritional value?** *Science* 2006, **311**:815-819.
  4. Benavente LM, Alonso JM: **Molecular mechanisms of ethylene signaling in *Arabidopsis*.** *Mol Biosys* 2006, **2**:165-173.
  5. Kendrick MD, Chang C: **Ethylene signaling: new levels of complexity and regulation.** *Curr Opin Plant Biol* 2008, **11**:479-485.
  6. Hahn A, Harter K: **Mitogen-Activated Protein kinase cascades and ethylene — signaling, biosynthesis or both?** *Plant Physiol* 2008, **149**:1207-1210.
  7. Zhong S, Lin Z, Grierson D: **Tomato ethylene receptor-CTR interactions: visualization of NEVER-RIPE interactions with multiple CTRs at the endoplasmic reticulum.** *J Exp Bot* 2008, **59**:965-972.
- This paper proposes a model whereby the NEVER-RIPE ethylene receptor interacts with several members of the CTR family and can potentially transmit information through a number of signaling pathways.
8. Barry CS, Giovannoni JJ: **Ripening in the tomato *Green-ripe* mutant is inhibited by ectopic expression of a protein that disrupts ethylene signaling.** *Proc Natl Acad Sci U S A* 2006, **103**:7923-7928.
  9. Resnick JS, Wen CK, Shockey JA, Chang C: **REVERSION-TO-ETHYLENE SENSITIVITY1, a conserved gene that regulates ethylene receptor function in *Arabidopsis*.** *Proc Natl Acad Sci U S A* 2006, **103**:7917-7922.
  10. Barry CS, McQuinn RP, Thompson AJ, Seymour GB, Grierson D, Giovannoni JJ: **Ethylene insensitivity conferred by the *Green-ripe* and *Never-ripe 2* ripening mutants of tomato.** *Plant Physiol* 2005, **138**:267-275.
  11. Kevany BM, Tieman DM, Taylor MG, Dal Cin V, Klee HJ: **Ethylene receptor degradation controls the timing of ripening in tomato fruit.** *Plant J* 2007, **51**:458-467.
  12. Lin Z, Arciga-Reyes L, Zhong S, Alexander L, Hackett R, Wilson I, Grierson D: **SITPR1, a tomato tetratricopeptide repeat protein, interacts with the ethylene receptors NR and LeETR1, modulating ethylene and auxin responses and development.** *J Exp Bot* 2008, **59**:4271-4287.
  13. Vrebalov J, Ruezinsky D, Padmanabhan V, White R, Medrano D, Drake R, Schuch W, Giovannoni J: **A MADS-box gene necessary for fruit ripening at the tomato *Ripening-inhibitor (Rin)* locus.** *Science* 2002, **296**:343-346.
  14. Manning K, Tor M, Poole M, Hong Y, Thompson AJ, King GJ, Giovannoni JJ, Seymour GB: **A naturally occurring epigenetic mutation in a gene encoding an SBP-box transcription factor inhibits tomato fruit ripening.** *Nat Genet* 2006, **38**:948-952.
  15. Hileman LC, Sundstrom JF, Litt A, Chen M, Shumba T, Irish VF: **Molecular and phylogenetic analyses of the MADS-Box gene family in tomato.** *Mol Biol Evol* 2006, **23**:2245-2258.
  16. Lin Z, Hong Y, Yin M, Li C, Zhang K, Grierson D: **A tomato HD-Zip homeobox protein, LeHB-1, plays an important role in floral organogenesis and ripening.** *Plant J* 2008, **55**:301-310.
- ACC oxidase has long been known to represent a key control point in ethylene biosynthesis. This important paper reports the identification of a homeobox protein that regulates ACC oxidase transcript levels, as well as the identification of associated regulatory elements in other ripening-associated genes, potentially placing it in a central location in ripening regulation.
17. Davuluri GR, van Tuinen A, Fraser PD, Manfredonia A, Newman R, Burgess D, Brummell DA, King SR, Palys J, Uhlig J et al.: **Fruit-specific RNAi-mediated suppression of *DET1* enhances carotenoid and flavonoid content in tomatoes.** *Nat Biotechnol* 2005, **23**:890-895.
  18. Alba R, Payton P, Fei ZJ, McQuinn R, Debbie P, Martin GB, Tanksley SD, Giovannoni JJ: **Transcriptome and selected metabolite analyses reveal multiple points of ethylene control during tomato fruit development.** *Plant Cell* 2005, **17**:2954-2965.
  19. Vicente AR, Saladié M, Rose JKC, Labavitch JM: **The linkage between cell wall metabolism and fruit softening: looking to the future.** *J Sci Food Agric* 2007, **87**:1435-1448.
  20. Goulao LF, Oliveira CM: **Cell wall modifications during fruit ripening: when a fruit is not the fruit.** *Trends Food Sci Technol* 2007, **19**:4-25.
  21. Guillon F, Philippe S, Bouchet B, Devaux MF, Frasse P, Jones B, Bouzayen M, Lahaye M: **Down-regulation of an auxin response factor in the tomato induces modification of fine pectin structure and tissue architecture.** *J Exp Bot* 2008, **59**:273-288.
  22. Faria MV, Maluf WR, Vilela de Resende JT, Andrade-Junior VC, do Nascimento IR, Gandolfi Benites FR, de Menezes CB,

- Azevedo SM: *Rin*, *nor<sup>A</sup>*, *og<sup>c</sup>* and *hp* mutants in tomatoes with different genetic backgrounds. *Pesquisa Agropecuaria Brasileira* 2006, **41**:793-800.
23. Ezura H, Owino WO: **Melon, an alternative model plant for elucidating fruit ripening.** *Plant Sci* 2008, **175**:121-129.
  24. Saladié M, Matas AJ, Isaacson T, Jenks MA, Goodwin SM, Niklas KJ, Xiaolin R, Labavitch JM, Shackel KA, Fernie AR *et al.*: **A reevaluation of the key factors that influence tomato fruit softening and integrity.** *Plant Physiol* 2007, **144**:1012-1028.  
This study emphasizes the idea that loss of fruit firmness, and consequently quality, is not solely dependent on cell wall metabolism and proposes that other factors, such as water status and cuticle structure should be examined carefully when considering targets for increasing shelf-life.
  25. Hovav R, Chehanovsky N, Moy M, Jetter R, Schaffer AA: **The identification of a gene (*Cwp1*), silenced during *Solanum* evolution, which causes cuticle microfissuring and dehydration when expressed in tomato fruit.** *Plant J* 2007, **52**:627-639.
  26. Leide J, Hildebrandt U, Reussing K, Riederer M, Vogg G: **The developmental pattern of tomato fruit wax accumulation and its impact on cuticular transpiration barrier properties: effects of a deficiency in a beta-ketoacyl-coenzyme A synthase (*LeCER6*).** *Plant Physiol* 2007, **144**:1667-1679.  
This paper highlights the importance of fruit waxes in limiting transpirational water, which in turn has important implications in identifying targets to improve shelf-life.
  27. Thomas TR, Shackel KA, Matthews MA: **Mesocarp cell turgor in *Vitis vinifera* L. berries throughout development and its relation to firmness, growth, and the onset of ripening.** *Planta* 2008, **228**:1067-1076.  
The significance of turgor pressure for fruit quality is not often discussed and this study of grape berry ripening provides convincing evidence that water status is a critical element in the equation.
  28. Wada H, Shackel KA, Matthews MA: **Fruit ripening in *Vitis vinifera*: apoplastic solute accumulation accounts for pre-veraison turgor loss in berries.** *Planta* 2008, **227**:1351-1361.
  29. Fraser PD, Enfissi EM, Bramley PM: **Genetic engineering of carotenoid formation in tomato fruit and the potential application of systems and synthetic biology approaches.** *Arch Biochem Biophys* 2008 (Epub ahead of print: doi:10.1016/j.abb.2008.10.009).
  30. Wang S, Liu J, Feng Y, Niu X, Giovannoni J, Liu Y: **Altered plastid levels and potential for improved fruit nutrient content by downregulation of the tomato DDB1-interacting protein CUL4.** *Plant J* 2008, **55**:89-103.
  31. Galpaz N, Wang Q, Menda N, Zamir D, Hirschberg J: **Abscisic acid deficiency in the tomato mutant *high-pigment 3* leading to increased plastid number and higher fruit lycopene content.** *Plant J* 2008, **53**:717-730.
  32. Fraser PD, Enfissi EM, Halket JM, Truesdale MR, Yu D, Gerrish C, Bramley PM: **Manipulation of phytoene levels in tomato fruit: effects on isoprenoids, plastids, and intermediary metabolism.** *Plant Cell* 2007, **19**:3194-3211.
  33. Giorio G, Stigliani AL, D'Ambrosio C: **Agronomic performance and transcriptional analysis of carotenoid biosynthesis in fruits of transgenic HighCaro and control tomato lines under field conditions.** *Transgenic Res* 2007, **16**:15-28.
  34. Butelli E, Titta L, Giorgio M, Mock HP, Matros A, Peterek S, Schijlen EG, Hall RD, Bovy AG, Luo J, Martin C: **Enrichment of tomato fruit with health-promoting anthocyanins by expression of select transcription factors.** *Nat Biotechnol* 2008, **26**:1301-1308.  
This is an excellent example of targeting a specific biosynthetic pathway, in this case that of tomato anthocyanins, to generate fruit with substantially increased nutritional value.
  35. Espley RV, Hellens RP, Putterill J, Stevenson DE, Kutty-Amma S, Allan AC: **Red colouration in apple fruit is due to the activity of the MYB transcription factor, MdMYB10.** *Plant J* 2007, **49**:414-427.
  36. Espley RV, Brendolise C, Chagne D, Kutty-Amma S, Green S, Volz R, Putterill J, Schouten HJ, Gardiner SE, Hellens RP, Allan AC: **Multiple repeats of a promoter segment causes transcription factor autoregulation in red apples.** *Plant Cell* 2009, tpc.108.059329.
  37. Schijlen EG, de Vos CH, Martens S, Jonker HH, Rosin FM, Molthoff JW, Tikunov YM, Angenent GC, van Tunen AJ, Bovy AG: **RNA interference silencing of chalcone synthase, the first step in the flavonoid biosynthesis pathway, leads to parthenocarpic tomato fruits.** *Plant Physiol* 2007, **144**:1520-1530.
  38. Bogs J, Jaffe FW, Takos AM, Walker AR, Robinson SP: **The grapevine transcription factor VvMYBPA1 regulates proanthocyanidin synthesis during fruit development.** *Plant Physiol* 2007, **143**:1347-1361.
  39. Luo J, Butelli E, Hill L, Parr A, Niggeweg R, Bailey P, Weisshaar B, Martin C: **AtMYB12 regulates caffeoyl quinic acid and flavonol synthesis in tomato: expression in fruit results in very high levels of both types of polyphenol.** *Plant J* 2008, **56**:316-326.
  40. Phillips PC: **Epistasis — the essential role of gene interactions in the structure and evolution of genetic systems.** *Nat Rev Genet* 2008, **9**:855-867.
  41. Causse M, Chaib J, Lecomte L, Buret M, Hospital F: **Both additivity and epistasis control the genetic variation for fruit quality traits in tomato.** *Theor Appl Genet* 2007, **115**:429-442.  
This study highlights the exciting potential of genetic approaches and the exploitation of existing germplasm, rather than targeting genetic engineering, to determine the molecular basis of complex fruit quality traits.
  42. Stevens R, Buret M, Duffe P, Garchery C, Baldet P, Rothan C, Causse M: **Candidate genes and quantitative trait loci affecting fruit ascorbic acid content in three tomato populations.** *Plant Physiol* 2007, **143**:1943-1953.
  43. Obando J, Fernandez-Trujillo JP, Martinez JA, Alarcon AL, Eduardo I, Arus P, Monforte AJ: **Identification of melon fruit quality quantitative trait loci using near-isogenic lines.** *J Am Soc Hort Sci* 2008, **133**:139-151.
  44. Cuevas HE, Staub JE, Simon PW, Zalapa JE, McCreight JD: **Mapping of genetic loci that regulate quantity of beta-carotene in fruit of US Western Shipping melon (*Cucumis melo* L.).** *Theor Appl Genet* 2008, **117**:1345-1359.
  45. Moreno E, Obando JM, Dos-Santos N, Fernandez-Trujillo JP, Monforte AJ, Garcia-Mas J: **Candidate genes and QTLs for fruit ripening and softening in melon.** *Theor Appl Genet* 2008, **116**:589-602.
  46. Gorguet B, Eggink PM, Ocana J, Tiwari A, Schipper D, Finkers R, Visser RG, van Heusden AW: **Mapping and characterization of novel parthenocarp QTLs in tomato.** *Theor Appl Genet* 2008, **116**:755-767.
  47. Park S, Cheng NH, Pittman JK, Yoo KS, Park J, Smith RH, Hirschi KD: **Increased calcium levels and prolonged shelf life in tomatoes expressing *Arabidopsis* H+.** *Plant Physiol* 2005, **139**:1194-1206.
  48. Morris J, Hawthorne KM, Hotze T, Abrams SA, Hirschi KD: **Nutritional impact of elevated calcium transport activity in carrots.** *Proc Natl Acad Sci U S A* 2008, **105**:1431-1435.
  49. Venneria E, Fanasca S, Monastra G, Finotti E, Ambra R, Azzini E, Durazzo A, Foddai MS, Maiani G: **Assessment of the nutritional values of genetically modified wheat, corn, and tomato crops.** *J Agric Food Chem* 2008, **56**:9206-9214.
  50. Deng P, Zhou X, Yang D, Hou H, Yang X, Li Y, Yang Y, Wang X, Fang S, Wu S, Liu J: **The definition, source, manifestation of and assessment of unintended effects in genetically modified plants.** *J Sci Food Agric* 2008, **88**:2401-2413.
  51. Finamore A, Roselli M, Britti S, Monastra G, Ambra R, Turrini A, Mengheri E: **Intestinal and peripheral immune response to MON810 maize ingestion in weaning and old mice.** *J Agric Food Chem* 2008, **56**:11533-11539.
  52. Hospital F: **Challenges for effective marker-assisted selection in plants.** *Genetica* 2008 (doi:10.1007/s10709-008-9307-1).
  53. Kevany BM, Taylor MG, Klee HJ: **Fruit-specific suppression of the ethylene receptor *LeETR4* results in early-ripening tomato fruit.** *Plant Biotechnol J* 2008, **6**:295-300.
  54. Shih CH, Chen Y, Wang M, Chu IK, Lo C: **Accumulation of isoflavone genistin in transgenic tomato plants overexpressing a soybean isoflavone synthase gene.** *J Agric Food Chem* 2008, **56**:5655-5661.